

Genuine Bayer® Aspirin Tablets

Drug Facts

Active ingredient (in each tablet)

Purposes

Aspirin 325 mg (NSAID)*......Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Hene

- temporarily relieves minor aches and pains due to:
 - headache
- muscle pain
- toothache
- menstrual pain
 colds
- minor pain of arthritis
- · temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

• hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a

prescription drug for

• gout • diabetes • arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint vomit blood have bloody or black stools
- have stomach pain that does not get better
 pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs



Genuine Bayer® Aspirin Tablets

If pregnant or breast-feeding, ask a health professional before use.
It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- · save carton for full directions and warnings
- Store at room temperature

Inactive ingredients corn starch, hypromellose, powdered cellulose, triacetin

Questions or comments? 1-800-331-4536 (Mon – Fri 9AM – 5PM EST)