



Midol® Complete Caffeine Free Caplets

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 500 mg.....	Pain reliever
Pamabrom 25 mg.....	Diuretic
Pyrilamine maleate 15 mg.....	Antihistamine

Uses for the temporary relief of these symptoms associated with menstrual periods:

- cramps
- bloating
- water-weight gain
- headache
- backache
- muscle aches

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



Midol® Complete Caffeine Free Caplets

Directions

- do not take more than the recommended dose
- adults and children 12 years and older:
 - take 2 caplets with water
 - repeat every 6 hours, as needed
 - do not exceed 6 caplets per day
- children under 12 years: consult a doctor

Other information Store at room temperature. Avoid temperatures above 25°C (77°F).

Inactive ingredients carnauba wax, croscarmellose sodium, D&C red #30 aluminum lake, D&C yellow #10 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, propylene glycol, shellac, titanium dioxide, triacetin

Questions or comments? 1-800-331-4536 (Mon-Fri 9AM - 5PM EST)