

Alka-Seltzer Plus® Maximum Strength Cold & Cough Liquid Gels

Drug Facts

Active ingredients (in each capsule) **Purposes** Chlorpheniramine maleate 2 mg......Antihistamine Dextromethorphan hydrobromide 10 mg.....Cough suppressant Phenylephrine hydrochloride 5 mg.....Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains
- nasal and sinus congestion
- sinus pressure
- cough
- · sore throat

headache

- runny nose
- sneezing
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:
- skin reddening blisters rash
 - hives
- facial swelling
 asthma (wheezing)
 shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver diseaseheart disease
- high blood pressure

- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- · persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- · do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- · be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children



Alka-Seltzer Plus® Maximum Strength Cold & Cough Liquid Gels

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.
 These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Other information

store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients D&C red #33, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)