



Coricidin[®] HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

Coricidin HBP[®] Maximum Strength Cold & Flu Day Liquid Gels

Drug Facts

Active ingredients (in each capsule)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Guaifenesin 200 mg.....	Expectorant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that is accompanied by excessive phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical



Coricidin® HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after taking the last Night dose before taking the Day product.
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 6 capsules in 12 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- store at 15°-25°C (59°-77°F)

Inactive ingredients FD&C yellow # 6, gelatin, glycerin, iron oxide black, lecithin, medium-chain triglycerides, mica, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)



Coricidin[®] HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

Coricidin[®] HBP Maximum Strength Cold & Flu Night Liquid Gels

Drug Facts

Active ingredients (in each capsule)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - cough
 - sore throat
 - runny nose
 - sneezing
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age



Coricidin[®] HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

Ask a doctor before use if you have

- liver disease
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after taking the last Day dose before taking the Night product.
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 4 capsules in 12 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- store at 15°-25°C (59°-77°F)

Inactive ingredients D&C yellow No.10, FD&C blue No.1, gelatin, glycerin, lecithin, medium-chain triglycerides, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, soybean oil, titanium dioxide

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)