



Coricidin® HBP

Maximum Strength Cold, Flu & Chest Congestion Liquid Gels

Drug Facts

Active ingredients (in each capsule)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Guaifenesin 200 mg.....	Expectorant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that is accompanied by excessive phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



Coricidin® HBP

Maximum Strength Cold, Flu & Chest Congestion Liquid Gels

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- store at 15°-25°C (59°-77°F)

Inactive ingredients FD&C yellow # 6, gelatin, glycerin, iron oxide black, lecithin, medium-chain triglycerides, mica, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)