



Coricidin HBP Maximum Strength Cold, Cough & Flu (Liquid)

Drug Facts

Active ingredients (in each 30 mL)

Purposes

Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg.....Cough suppressant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - cough
 - sore throat
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that is accompanied by excessive phlegm (mucus)
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.



Coricidin HBP Maximum Strength Cold, Cough & Flu (Liquid)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than recommended dose
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 4 hours. Do not exceed 150 mL (5 doses) in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- **each 30 mL contains:** sodium 38 mg
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red # 40, flavor, glycerin, maltitol solution, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-317-2165 (Mon – Fri 9AM – 5PM EST)