



# Phillips' Stool Softener Liquid Gels

## Drug Facts

<b>Active ingredient (in each softgel)</b>	<b>Purpose</b>
Docusate sodium 100 mg.....	Stool softener laxative

### Use

- for relief of occasional constipation (irregularity)
- this product generally produces a bowel movement in 12 to 72 hours

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if you have

- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

**Ask a doctor or pharmacist before use if you are** presently taking mineral oil

#### Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- take softgels with a full glass (8 oz) of water

adults and children 12 years and older	take 1 to 3 softgels daily or as directed by a doctor. This dose may be taken as a single daily dose or in divided doses.
children 6 to under 12 years	take 1 softgel daily or as directed by a doctor
children under 6 years of age	ask a doctor

### Other information

- **each softgel contains:** sodium 6 mg
- very low sodium
- store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive ingredients** FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

**Questions or comments?** 1-800-986-0369

(Mon – Fri 9AM – 5PM EST)