



Midol Long Lasting Relief Caplets

Drug Facts	
Active ingredient (in each caplet)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Uses	
<ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ muscular aches ■ backache ■ minor pain of arthritis ■ toothache ■ premenstrual and menstrual cramps ■ headache ■ the common cold ■ temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> ■ more than 6 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ skin reddening ■ blisters ■ rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p>	
Do not use	
<ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product 	
Ask a doctor before use if you have liver disease	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present <p>These could be signs of a serious condition.</p>	
<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	
Directions ■ do not take more than directed (see overdose warning)	
adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	■ do not use



Midol Long Lasting Relief Caplets

Other information

- store at 20-25°C (68-77°F)
- **do not use if printed foil under cap is broken or missing**
- meets the requirements of USP *Dissolution Test 4*

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments? 1-800-331-4536 (Mon - Fri 9AM – 5PM EST)